

UNSAFE	SAFE
<p>Avoid pain medications that contain Opiates(e.g. Codein):</p> <ul style="list-style-type: none"> • Tylenol 1,2,3, or 4(all Opioids) • Demerol • Percet • Fiorinal Plan ¼ or ½ • Levo-Dromoran • 222,282,292,692,Darvon(Propoxyphene) • Talwin • Gabapentin - lowest Dosage 300 mg. • Perodan • Leritine • Dilaudid • Nabilone <p>Avoid Nerve and Sleeping Pills including:</p> <ul style="list-style-type: none"> • Librium • Valium • Clonazepam *All Med with "pam" suffix will most likely • Tranxene be in same group. • Serax • Xanax • Diazepan <p>Avoid Sleeping Pills including these and others:</p> <ul style="list-style-type: none"> • Dalmane • Halcion • Restoril • Tuinal • Seconal • Zopiclone(Imovane) <p>Ovoid Muscle Relaxants:</p> <ul style="list-style-type: none"> • Robaxisal • Robaxacet • Parafon • Flexeril • Methocarbamol • Cyobenzapaprine <p>Over the Counter Medications can be a Serious Threat:</p> <ul style="list-style-type: none"> • Cough syrups contain alcohol, codeine and antihistamines. These are all drugs which need to be avoided. <p>Avoid Sedating Antihistamines such as:</p> <ul style="list-style-type: none"> • Gravol • Mimenhydriate • Actifed • Dimetap • Chlortripion • Benydryl or products containing 	<p>Pain Medications:</p> <ul style="list-style-type: none"> • Regular or Extra Strength Tylenol • ASA or Aspirin • Advil or Ibuprofen • Midol <p>Available only by Prescription:</p> <ul style="list-style-type: none"> • Tryptan • Buspirone(Buspar) • Gapapentin • Toradol • Possible other prescription medications-please contact Resident Nurse for clarification. <p>Antidepressants Safe with Proper Use and by Prescription Only:</p> <ul style="list-style-type: none"> • Elavil • Citalopram • Morex • Serzone • Desipramine • Effexor(Venlafaxine) • Zoloft(Sertraline) • Prozac(Fluxentine) • Luvox(Fluvoxamine) • Paxil(Paroxetine) • Trazodone(Desyrel) • Mirtazapine • Bupropion • Seroquel(Quetiapine) <p>Migraines:</p> <ul style="list-style-type: none"> • Lmitrex <p>Non-Sedating Antihistamines:</p> <ul style="list-style-type: none"> • Seldane • Claritin • Hismanil <p>Sleep Aids:</p> <p>Epsom Salt</p> <ul style="list-style-type: none"> • Melatonin • Calcium(33 mg) Magnesium(167mg) with VD3(5mcg) • Lavender Oil

Note:This is a partial list. If you require more information, please ask the Doctor or Pharmacist about non-psycho active mood altering Medications brought into treatment and taken in the two weeks prior to Intake date with result in the Client's immediate discharge from the program.